

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

4


## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6:  C07C 275/28, C07D 213/75, 257/04, 231/12, 401/12, A61K 31/17, 31/44, 31/41, C07D 213/40, 307/38, 277/28, 233/54, C07C 311/21, C07D 333/20		A1	(11) International Publication Number: <b>WO 96/10559</b>  (43) International Publication Date: 11 April 1996 (11.04.96)
(21) International Application Number: PCT/JP95/01982  (22) International Filing Date: 29 September 1995 (29.09.95)		(74) Agent: SEKI, Hideo; Fujisawa Pharmaceutical Co., Ltd., Osaka Factory, 1-6, Kashima 2-chome, Yodogawa-ku, Osaka-shi, Osaka 532 (JP).	
(30) Priority Data:  9419970.0 4 October 1994 (04.10.94) GB 9506720.3 31 March 1995 (31.03.95) GB 9514021.6 10 July 1995 (10.07.95) GB		(81) Designated States: AU, CA, CN, HU, JP, KR, MX, RU, US. European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).	
(71) Applicant (for all designated States except US): FUJISAWA PHARMACEUTICAL CO., LTD. [JP/JP]; 4-7, Doshimachi 3-chome, Chuo-ku, Osaka-shi, Osaka 541 (JP).  (72) Inventors; and (75) Inventors/Applicants (for US only): TERASAWA, Takeshi [JP/JP]; 1625-302, Matsugakcanakamachi, Kawachinagano-shi, Osaka 586 (JP). TANAKA, Akira [JP/JP]; 9-10-302, Nakano-cho, Takamizuka-shi, Hyogo 665 (JP). CHIBA, Toshiyuki [JP/JP]; 1-1-503, Nakatsuji-cho, Nara-shi, Nara 630 (JP). TAKASUGI, Hisashi [JP/JP]; 3-116-10, Mazu Umekita, Sakai-shi, Osaka 591 (JP).		Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.	

## (54) Title: UREA DERIVATIVES AND THEIR USE AS ACAT-INHIBITORS

## (57) Abstract

Urea derivatives of formula (I), wherein R<sup>1</sup> is a group of formula (I) (in which R<sup>4</sup> is aryl which may have suitable substituent(s), or heterocyclic group which may have suitable substituent(s), and Y is bond, lower alkylene, -S-, -O-, (a), -CH-, -CONH-, (b), (in which R<sup>7</sup> is lower alkyl), -NHSO<sub>2</sub>-, -SO<sub>2</sub>NH-, -SO<sub>2</sub>NHCO- or -CONHSO<sub>2</sub>-); or thiazolyl, imidazolyl, pyrazolyl, pyridyl, thienyl, furyl, isoxazolyl or chromanyl, each of which may have suitable substituent(s); R<sup>2</sup> is lower alkyl, lower alkoxy(lower)alkyl, cycloalkyl, ar(lower)alkyl which may have suitable substituent(s), heterocyclic group or heterocyclic(lower)alkyl, R<sup>3</sup> is aryl which may have suitable substituent(s) or heterocyclic group which may have suitable substituent(s), and n is 0 or 1, and a pharmaceutically acceptable salt thereof which are useful as a medicament in the treatment of hypercholesterolemia, hyperlipidemia and atherosclerosis.




WO 96/10559

PCT/JP95/01982


- 210 -

## C L A I M S

## 1. A compound of the formula :



wherein

 $R^1$  is a group of the formula :

(in which

$R^4$  is aryl which may have suitable substituent(s), or heterocyclic group which may have suitable substituent(s), and

$Y$  is bond, lower alkylene,  $-S-$ ,  $-O-$ ,  $-C(=O)-$ ,  
 $=CH-$ ,  $-CONH-$ ,  $-N(CO)-$ , (in which  $R^7$  is lower alkyl),  
 $-NHSO_2-$ ,  $-SO_2NH-$ ,  $-SO_2NHCO-$  or  $-CONHSO_2-$ );  
 or

thiazolyl, imidazolyl, pyrazolyl, pyridyl,  
 thiienyl, furyl, isoxazolyl or chromanyl, each of  
 which may have suitable substituent(s);

$R^2$  is lower alkyl, lower alkoxy(lower)alkyl,  
 cycloalkyl, ar(lower)alkyl which may have  
 suitable substituent(s), heterocyclic group or  
 heterocyclic(lower)alkyl,

$R^3$  is aryl which may have suitable substituent(s) or  
 heterocyclic group which may have suitable

WO 96/10559


PCT/JP95/01982

- 211 -

substituent(s), and  
 n is 0 or 1,  
 and a pharmaceutically acceptable salt thereof.

5 2. A compound of claim 1, wherein  
 R<sup>1</sup> is a group of the formula :

10.



(in which  
 15 R<sup>4</sup> is phenyl which may have 1 to 3 substituent(s)  
 selected from the group consisting of  
 halogen, lower alkyl, di(lower)alkylamino,  
 protected amino, cyano, heterocyclic group  
 which may have mono(or di or tri)-  
 ar(lower)alkyl, hydroxy, protected hydroxy  
 and mono(or di or tri)halo(lower)alkyl;  
 20 or thienyl, pyrazolyl, imidazolyl,  
 triazolyl, pyridyl, pyrrolyl, tetrazolyl,  
 oxazolyl, thiazolyl, oxadiazolyl,  
 piperazinyl, thiazolidinyl or  
 methylenedioxyphenyl, each of which may have  
 25 1 to 3 substituent(s) selected from the  
 group consisting of lower alkyl, mono(or di  
 or tri)ar(lower)alkyl and oxo;

30 Y is bond, lower alkylene, -S-, -O-, -C-, =CH-,  
 -CONH-, -N-CO- (in which R<sup>7</sup> is lower alkyl),  
 R<sup>7</sup>  
 -NHSO<sub>2</sub>-, -SO<sub>2</sub>NH-, -SO<sub>2</sub>NHCO- or -CONHSO<sub>2</sub>-);  
 or  
 35 thiazolyl, imidazolyl, pyrazolyl, pyridyl,